Computer Algorithms Support Physician Decisions in Traumatic Head Injury

Caroline L.S. George, MD, Nancy S. Harper, MD

Do not let the title of the article in this issue of *Pediatrics*, "Traumatic Head Injury and the Diagnosis of Abuse: A Cluster Analysis," by Boos et al[1] intimidate those of you who are not biostatisticians. This article is important for providers who evaluate children with head injuries in the hospital setting. Young children remain disproportionately affected by child maltreatment, with the majority of fatalities occurring as a result of abusive head trauma (AHT).[2] There has been extensive, high-quality research into clinical and radiologic findings associated with AHT,[3–5] and multicenter research networks have been formed to accelerate this important work. One of the earliest networks, the Pediatric Brain Injury Research Network (PediBIRN), has prospectively collected data on children <3 years of age with symptomatic head injury in the PICU setting to identify a valid and reliable clinical prediction rule (CPR) for pediatric AHT.[6]

The authors present a unique and important analysis of the clinical and radiologic findings of 500 young patients with acute neurotrauma from data collected by PediBIRN.[7] Their hypothesis is that well-established computer algorithms used to sort "like" data would identify subgroups of children with discreet clinical findings. Furthermore, these subgroups would have specific relationships to indicators of abuse and physician diagnoses. Why use computer algorithms when clinical variables specific to AHT are available? The application of computer algorithms to clinical data broadly characterizes these mathematical clustering methods with the goal of identifying more subtle associations less apparent to researchers and clinicians. There are examples of this approach in the medical sciences, such as identifying phenotypically different asthma groups[8] or recognizing distinct clinical subtypes of Alzheimer disease.[9] The use of computer algorithms makes this research unique, allowing elimination of potential researcher bias as the computer is blinded to physician diagnoses and previously published associations.

Several key points are integral in understanding the importance of the authors' cluster analysis. First, they used an "unsupervised" cluster analysis, meaning that computer algorithms analyzed unlabeled data and sorted data into groups without a target outcome. Next, the authors used well-established cluster modeling tools.[9,10] K-means is a partitioning cluster algorithm tool where each cluster has a center (eg, the color orange might be in a cluster with yellow and red but would not be in a cluster centered around blue). Divisive and agglomerative algorithms are hierarchical tools that either start with one larger cluster and divide like items into smaller groups or start with smaller groups and connect them back to a larger core group, respectively.

Department of Pediatrics, University of Minnesota, University of Minnesota Masonic Children's Hospital, Minneapolis, Minnesota

DOI: https://doi.org/10.1542/peds.2021-054009

Accepted for publication Oct 20, 2021

Address correspondence to Caroline L.S. George, MD, *Otto Bremer Trust* Center for Safe and Healthy Children, University of Minnesota, 2512 S 7th St, Suite R107, Minneapolis, MN 55454. E-mail: cgeorge@umn.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2022 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: Drs George and Harper's institution has received payment for expert witness testimony that has been provided in cases of suspected child abuse in which the physicians have been subpoenaed to testify.

COMPANION PAPER: A companion to this article can be found online at www.pediatrics.org/cgi/doi/10.1542/peds.2021-051742.

To cite: George CL.S, Harper NS. Computer Algorithms Support Physician Decisions in Traumatic Head Injury. *Pediatrics*. 2022;149(1):e2021054009

Downloaded from http://publications.aap.org/pediatrics/article-pdf/149/1/e2021054009/1226914/peds_2021054009.pdf by UT Rio Grande Valley user on 28 January 2022

For example, a phylogenetic tree of birds would demonstrate a closer ancestor between a falcon and an eagle rather than an ostrich.[11] Each tool has strengths and weaknesses; therefore, several tools are used to sort data and analyze cluster similarities and differences. Finally, the authors assessed data partitioned by each algorithm with the "triad," defined as the presence of 3 clinical variables: any subdural hemorrhage (SDH); any encephalopathy before presentation for care; and retinal hemorrhages (RHs) described by an ophthalmologist as dense, extensive, covering a large surface area, and/or extending to the ora serrata. The odds ratio (OR) and confidence interval are presented with respect to substantial association with high likelihood of AHT (HiAHT) (OR >10) and low likelihood of AHT (LoAHT) (OR <0.01). Many clinical findings did not strongly partition with either group but still demonstrated statistical significance.

The cluster-modeling tools optimally partitioned the data into 2 groups, one group comprised the majority of children ultimately diagnosed by a physician with AHT (therefore HiAHT). Variables that had the most significant association with the HiAHT group included signs/symptoms of significant brain dysfunction: brain hypoxemia, ischemia, and/or swelling; acute encephalopathy (persistent); respiratory compromise prior to admission (PTA); presence of SDH or fluid collection; and retinoschisis. Not surprisingly, acute encephalopathy, SDH, and extensive retinal hemorrhages (RHs) also had significant associations with triad partitioning. However, it is interesting to see that the triad, as defined by the authors, was not as sensitive a partitioning tool as the other cluster tools used in this study. Also of interest, additional injuries seen in child physical abuse, such as bruising (ear, neck, torso) and fractures (considered moderately/highly specific for abuse), were statistically significant in the cluster analysis, but did not have substantial associations with HiAHT or LoAHT.

What does this complex analysis mean? Children admitted with symptomatic traumatic brain injury (TBI) are not on a continuum in which the presentation with AHT looks similar to accidental head trauma or LoAHT. Presentations strongly associated with AHT are characterized by CNS dysfunction, including respiratory compromise PTA and persistent encephalopathy, as well as radiologic evidence of substantial brain injury and SDH that would not be described as "contact" (unilateral). Additionally, children with AHT are more likely to have extensive RHs and/or retinoschisis, fractures, bruising, and/or abdominal injury, consistent with the published literature.[12] Presentations that are lower risk for AHT, seen more often with accidental head trauma, often have contact injuries seen with impact to the head, such as a skull fracture with an associated epidural or subdural hemorrhage. These associations are in line with published literature on objective clinical findings that assist in discriminating between these etiologies.[3–5]

A single clinical finding in isolation may not discriminate between children at high or low risk for AHT. However, this analysis and the current literature illustrate the predictive probability of clinical findings seen in combination.[13,14] The results of this unique cluster analysis, demonstrating the association of clinical findings of significant brain dysfunction with AHT, will assist medical providers in the hospital setting, highlighting the importance of a comprehensive medical evaluation in children with traumatic head injury.

ABBREVIATIONS

AHT: abusive head trauma
CNS: central nervous system
OR: odds ratio
PediBIRN: Pediatric Brain Injury Research Network
PTA: prior to admission
RH: retinal hemorrhage
SDH: subdural hemorrhage
TBI: traumatic brain injury

REFERENCES

1. Boos SC, Wang M, Karst WA, Hymel KP; Pediatric Brain Injury Research Network (PediBIRN) Investigators. Traumatic head injury and the diagnosis of abuse: a cluster analysis. *Pediatrics*. 2022:149(1):e2021051742
2. Palusci VJ, Covington TM. Child maltreatment deaths in the US National Child Death Review Case Reporting System. *Child Abuse Negl*. 2014;38(1):25–36
3. Maguire S, Pickerd N, Farewell D, Mann M, Tempest V, Kemp AM. Which clinical features distinguish inflicted from non-inflicted brain injury? A systematic review. *Arch Dis Child*. 2009;94(11):860–867
4. Kemp AM, Jaspan T, Griffiths J, et al. Neuroimaging: what neuroradiological features distinguish abusive from non-abusive head trauma? A systematic review. *Arch Dis Child*. 2011;96(12):1103–1112
5. Kelly P, John S, Vincent AL, Reed P. Abusive head trauma and accidental head injury: a 20-year comparative study of referrals to a hospital child protection team. *Arch Dis Child*. 2015;100(12):1123–1130
6. Hymel KP, Willson DF, Boos SC, et al; Pediatric Brain Injury Research Network (PediBIRN) Investigators. Derivation of a clinical prediction rule for pediatric abusive head trauma. *Pediatr Crit Care Med*. 2013;14(2):210–220
7. Pediatric Brain Injury Research Network. Welcome to the CDR Implementation Trial.

Downloaded from http://publications.aap.org/pediatrics/article-pdf/149/1/e2021054009/1226914/peds_2021054009.pdf by UT Rio Grande Valley user on 28 January 2022

Available at: https://www.pedibirn.com. Accessed October 3, 2021

8. Haldar P, Pavord ID, Shaw DE, et al. Cluster analysis and clinical asthma phenotypes. *Am J Respir Crit Care Med*. 2008;178(3):218–224
9. Alashwal H, El Halaby M, Crouse JJ, Abdalla A, Moustafa AA. The application of unsupervised clustering methods to Alzheimer's disease. *Front Comput Neurosci*. 2019;13:31
10. Everitt BS, Landau S, Leese M, Stahl D. *Cluster Analysis*, 5th ed. Chichester, England: John Wiley & Sons; 2011
11. Mayr GG. The deep divergences of neornithine birds: a phylogenetic analysis of morphological characters. *Cladistics*. 2003;19(6):527–533
12. Choudhary AK, Dies Suarez P, Binenbaum G, et al. Consensus statement on abusive head trauma: additional endorsements. *Pediatr Radiol*. 2019;49(3):421
13. Maguire SA, Kemp AM, Lumb RC, Farewell DM. Estimating the probability of abusive head trauma: a pooled analysis. *Pediatrics*. 2011;128(3):e550–e564
14. Vinchon M, de Foort-Dhellemmes S, Desurmont M, Delestret I. Confessed abuse versus witnessed accidents in infants: comparison of clinical, radiological, and ophthalmological data in corroborated cases. *Childs Nerv Syst*. 2010;26(5):637–645

Downloaded from http://publications.aap.org/pediatrics/article-pdf/149/1/e2021054009/1226914/peds_2021054009.pdf by UT Rio Grande Valley user on 28 January 2022